

UTAS-PHYS-02-09

hep-th/0209233

July 2002

Three-body phase space: symmetrical treatments*

A.I. Davydychev[†] and R. Delbourgo[‡]

*^aSchool of Mathematics and Physics, University of Tasmania,
GPO Box 252-21, Hobart, Tasmania 7001, Australia*

Abstract

We derive expressions for three-body phase space that are explicitly symmetrical in the masses of the three particles, by three separate methods.

*Presented by R. Delbourgo at the 15th Biennial Congress of the Australian Institute of Physics (Sydney, Australia, July 2002)

[†] On leave from Institute for Nuclear Physics, Moscow State University, 119899 Moscow, Russia.
Email address: davyd@thep.physik.uni-mainz.de

[‡]Email address: bob.delbourgo@utas.edu.au

Nature of the problem

The phase volume in D -dimensional space-time for the decay process $p \rightarrow 1 + 2 + 3$ can be reduced to the integral

$$\rho_{p \rightarrow 1+2+3} = \frac{(4\pi)^{1-D}(p^2)^{1-D/2}}{2\Gamma(D-2)} \iiint ds dt du \delta(s+t+u-m_1^2-m_2^2-m_3^2-p^2) [\Phi(s, t, u)]^{D/2-2} \theta(\Phi),$$

where $s = (p_1 + p_2)^2$, $t = (p_2 + p_3)^2$, $u = (p_3 + p_1)^2$ are the three Mandelstam variables and, in the physical region, the Kibble cubic

$$\begin{aligned} \Phi(s, t, u) = & stu - s(m_1^2 m_2^2 + p^2 m_3^2) - t(m_2^2 m_3^2 + p^2 m_1^2) - u(m_3^2 m_1^2 + p^2 m_2^2) \\ & + 2(m_1^2 m_2^2 m_3^2 + p^2 m_1^2 m_2^2 + p^2 m_2^2 m_3^2 + p^2 m_3^2 m_1^2). \end{aligned}$$

stays positive. Clearly ρ is symmetric in the three masses. However, when one eliminates one of the Mandelstam variables (or any linear combination) one is left, for even D , with an elliptic integral which is not *explicitly* symmetrical, although it must be so implicitly. The aim of this article is to exhibit three routes for getting a satisfyingly symmetrical result in the $4D$ case, where ρ is nothing but the area of the Dalitz-Kibble plot [1, 2] divided by $128\pi^3 p^2$. Detailed derivations are not given, for lack of space, but will be published elsewhere.

Route 1: Elliptic Drive

The complicated expressions found by Almgren [3] and Bauberger et al [4], obtained by finally integrating [$s_{1,2} = (m_1 \pm m_2)^2$, $s_{3,4} = (p \mp m_3)^2$ below]

$$\rho = \frac{1}{128\pi^3 p^2} \int_{s_2}^{s_3} \frac{ds}{s} \sqrt{(s-s_1)(s-s_2)(s_3-s)(s_4-s)},$$

involve the complete elliptic integrals, $K(k)$, $E(k)$ and $\Pi(\alpha^2, k)$, where

$$k = \sqrt{\frac{q_{+-}q_{-+}}{q_{++}q_{--}}} \quad \text{and} \quad q_{\pm\pm} = (p \pm m_2)^2 - (m_1 \pm m_2)^2.$$

The main point is that the result is symmetric under $1 \leftrightarrow 2$ interchange but *not obviously* so under $1 \leftrightarrow 3$ and $2 \leftrightarrow 3$ interchanges. The key to converting these expressions into symmetric form is to pass to the Jacobian zeta-functions Z via

$$\frac{\Pi(\alpha^2, k)}{K(k)} = 1 + \frac{\alpha Z(\beta, k)}{\sqrt{(1-\alpha^2)(k^2-\alpha^2)}}$$

with $\sin \beta = \alpha/k$ and to use the addition formula (142.01) of Byrd and Friedman [5]:

$$\begin{aligned} Z(\beta_1, k) \pm Z(\beta_2, k) &= Z(\varphi_{\pm}, k) \pm k^2 \sin \beta_1 \sin \beta_2 \sin \varphi_{\pm}, \\ \tan \frac{\varphi_{\pm}}{2} &= \frac{\sin \beta_1 \sqrt{1 - k^2 \sin^2 \beta_2} \pm \sin \beta_2 \sqrt{1 - k^2 \sin^2 \beta_1}}{\cos \beta_1 + \cos \beta_2}. \end{aligned}$$

After numerous manoeuvres, including identification of the three angles

$$\varphi_2 = \varphi_+, \quad \varphi_1 = \varphi_-, \quad \varphi_3 = \arctan \left[\frac{\sqrt{q_{++}q_{--}}}{2(pm_3 - m_1m_2)} \right],$$

etc. and noting the remarkable connection

$$Z(\varphi_1, k) + Z(\varphi_2, k) + Z(\varphi_3, k) = k^2 \sin \varphi_1 \sin \varphi_2 \sin \varphi_3,$$

one obtains the beautifully symmetric form:

$$\begin{aligned} \rho = \frac{\sqrt{q_{++}q_{--}}}{128\pi^3 p^2} & \left\{ (p^2 + m_1^2 + m_2^2 + m_3^2) [E(k) - K(k)] \right. \\ & \left. + \sqrt{q_{++}q_{--}} K(k) \left[\frac{Z(\varphi_1, k)}{\sin^2 \varphi_1} + \frac{Z(\varphi_2, k)}{\sin^2 \varphi_2} + \frac{Z(\varphi_3, k)}{\sin^2 \varphi_3} \right] \right\}. \end{aligned}$$

Route 2: Phase Cut Way

In the triangular phase plot, pictured in Figure 1, points O refer to configurations where the Mandelstam variables assume their minimum values, points P where they attain their maximum values and the Q -lines and R -lines are tangents at P and O . Needing to integrate within the cubic outline $OPOPOP$, one is guaranteed to obtain a symmetric answer by finding the area of the hexagon A_Q with Q vertices and cutting out the six OQP crescents. Now area

$$A_Q = \frac{1}{2} (p - m_1 - m_2 - m_3)^2 \left[(p + m_1 + m_2 + m_3)^2 - 4(m_1^2 + m_2^2 + m_3^2) \right],$$

while a typical crescent has an area

$$\begin{aligned} A_{O_t P_s Q_{su}} = \int_{s_{O_t}}^{(p-m_3)^2} \frac{ds}{2s} & \left[(p^2 - m_3^2)(m_2^2 - m_1^2) - s \left(s - p^2 - m_1^2 + m_2^2 + m_3^2 + 4m_2m_3 \right) \right. \\ & \left. - \sqrt{(s - s_1)(s - s_2)(s_3 - s)(s_4 - s)} \right] \end{aligned}$$

that is definitely given by an elliptic function. These crescents are readily evaluated in the limit of large p and of small $Q = p - (m_1 + m_2 + m_3)$ by analytical means. One finds to leading order

$$\begin{aligned} A_{O_t P_s Q_{su}} &= m_3^2 p^2 \left[\log \left(\frac{p}{m_2 + m_3} \right) - \frac{3}{2} \right] \\ &\quad - \frac{m_2 m_3^2 p^2}{12(m_2 + m_3)^4} \left[12m_3^3 + 23m_2 m_3^2 + 20m_2^2 m_3 + 6m_2^3 \right] + \mathcal{O}(p), \\ A_{O_t P_s Q_{su}} &= 2Q^2 \left[m_1 m_2 - \sqrt{m_1 m_2 m_3 (m_1 + m_2 + m_3)} \theta_{13} \right] + \mathcal{O}(Q^3), \\ \sin \theta_{13} &= \sqrt{\frac{m_1 m_3}{(m_1 + m_2)(m_2 + m_3)}}, \quad \text{etc.} \end{aligned}$$

$$i\Delta_c(x|m_i) = \frac{1}{(2\pi)^{D/2}} \left(\frac{m}{r}\right)^{D/2-1} K_{D/2-1}(mr) , \quad r = \sqrt{-x^2 + i\epsilon} .$$

The Fourier integral leads to an integral of a Bessel function J of the first kind with a product of three K functions of the second kind; balanced progress may be achieved via the expansion,

$$K_\nu(z) = \sqrt{\frac{\pi}{2z}} \left[1 + \frac{4\nu^2 - 1}{8z} + \dots \right] ,$$

which terminates for odd D but is asymptotic for even D . Since for example,

$$K_1(m_1 r) K_1(m_2 r) K_1(m_3 r) = \frac{\pi^{3/2} \exp[-(m_1 + m_2 + m_3)r]}{(8m_1 m_2 m_3 r^3)^{1/2}} \left[1 + \frac{3}{8r} \left(\frac{1}{m_1} + \frac{1}{m_2} + \frac{1}{m_3} \right) + \dots \right]$$

we are left with a doable integral, that produces a series of hypergeometric functions, whose leading term is

$$I_3(p) \sim \frac{1}{(\sigma\omega)^{(3-D)/2}} \Gamma\left(\frac{3-D}{2}\right) F\left(\frac{3-D}{4}, \frac{5-D}{4}; \frac{D}{2}; \frac{p^2}{\sigma^2}\right) ,$$

with $\sigma = m_1 + m_2 + m_3$, $\omega = m_1 m_2 m_3$. As imaginary parts of hypergeometric functions are [8]

$$\text{Im}F(a, b; c; z) = -\frac{\pi\Gamma(c)(z-1)^{c-a-b}\theta(z-1)}{\Gamma(a)\Gamma(b)\Gamma(1-a-b+c)} F(c-a, c-b; c-a-a+1; 1-z) ,$$

we deduce that

$$\rho \propto \frac{(p^2 - \sigma^2)^2}{\omega^{-1/2}\sigma^{7/2}} F\left(\frac{9}{4}, \frac{7}{4}; 3; 1 - \frac{p^2}{\sigma^2}\right) + \mathcal{O}\left((p^2 - \sigma^2)^3\right) ,$$

$$(p^2 - \sigma^2) = Q(2m_1 + 2m_2 + 2m_3 + Q) ,$$

and recognize the threshold behaviour found by other routes. This is a particular case of the general N -body threshold behaviour, $\rho_N(p) \sim Q^{(N-1)D/2-(N+1)/2}$. One last comment: it is best to input the massless propagators $i\Delta_c(x|0) = \Gamma(D/2-1)/(4\pi^{D/2}r^{D-2})$ at the stage of Fourier integration, rather than attempt to take the zero-mass limit of the above massive formulae, because that procedure is extremely tricky.

Acknowledgements

We are pleased to acknowledge support from the Australian Research Council under grant number A00000780.

References

- [1] R.H. Dalitz, Phil. Mag. **44** (1953) 1068.
- [2] T.W.B. Kibble, Phys. Rev. **117** (1960) 1159.
- [3] B. Almgren, Arkiv för Physik **38** (1968) 161.
- [4] S. Bauberger, F. Berends, M. Böhm and M. Buza, Nucl. Phys. **B434** (1995) 383.
- [5] P.F. Byrd and M.D. Friedman, *Handbook of Elliptic Integrals for Engineers and Physicists*, Springer, 1954.
- [6] S. Groote, J.G. Körner and A.A. Pivovarov, Nucl. Phys. **B542** (1999) 515.
- [7] S. Groote and A.A. Pivovarov, Nucl. Phys. **B580** (2000) 459.
- [8] A. Bashir, R. Delbourgo and M.L. Roberts, J. Math. Phys. 42 (2001) 5553.
- [9] A.I. Davydychev and V.A. Smirnov, Nucl. Phys. **B554** (1999) 391.